

University of Groningen

Fever-Induced Paroxysmal Weakness and Encephalopathy (FIPWE)-Part of a Phenotypic Continuum in Patients With ATP1A3 Mutations?

Sival, Deborah A.; Vansenne, Fleur; Van der Hout, Annemieke H.; Tijssen, Marina A. J.; de Koning, Tom J.

Published in:
Pediatric neurology

DOI:
[10.1016/j.pediatrneurol.2017.12.009](https://doi.org/10.1016/j.pediatrneurol.2017.12.009)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sival, D. A., Vansenne, F., Van der Hout, A. H., Tijssen, M. A. J., & de Koning, T. J. (2018). Fever-Induced Paroxysmal Weakness and Encephalopathy (FIPWE)-Part of a Phenotypic Continuum in Patients With ATP1A3 Mutations? *Pediatric neurology*, 81, 57-58. <https://doi.org/10.1016/j.pediatrneurol.2017.12.009>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Correspondence

Fever-Induced Paroxysmal Weakness and Encephalopathy (FIPWE)—Part of a Phenotypic Continuum in Patients With *ATP1A3* Mutations?



We read with interest the article by Yano et al.¹ identifying a new phenotype of patients with a heterozygous mutation in the gene encoding the sodium-potassium ATPase alpha-3 subunit (*ATP1A3*) at c.2267G>A (residue 756). Previously, specific *ATP1A3* gene mutations were associated with unique phenotypes, including alternating hemiplegia of childhood (AHC), rapid-onset dystonia-parkinsonism (RDP/DYT12), and cerebellar ataxia, pes cavus, optic atrophy, sensorineural hearing loss.² In children with *ATP1A3* gene mutations, Yano et al. described a group of patients with another phenotype, presenting with fever-induced paroxysmal muscle weakness and encephalopathy (FIPWE). The authors allocate this specific clinical presentation to a mutation at a single residue 756 (p.(Arg756Leu), p.(Arg756His), or p.(Arg756Cys)).¹

In addition to recognition of separate genotype-phenotype relationships, awareness of the phenotypic continuum of *ATP1A3* mutations is necessary.³ This is illustrated by three of our patients with FIPWE-like features, who were reported with different *de novo ATP1A3* gene mutations. The first two (unrelated) children were both diagnosed with an identical *de novo ATP1A3* gene mutation at c.2266C>T, p.(Arg756Cys) (NM_152296.4). These two children suffered from fever-induced attacks with paroxysmal muscle weakness, encephalopathy, ataxia, chorea, dystonia, and mutism. The third child was diagnosed with a *de novo ATP1A3* gene mutation at c. 2232C>A, p.(Asn744Lys) (NM_152296.4). He suffered from three severe fever- and stress-induced attacks with paroxysmal muscle weakness, asymmetrical dystonia, mutism, electroencephalographic seizure activity, and encephalopathy. All three children recovered in between attacks, albeit incomplete and slowly. In the first two children, the FIPWE phenotype-genotype relationship was established. The symptoms of the third patient, however, were attributed to a spectrum of overlapping acronyms (AHC, RPD, and FIPWE).

Although genetically different, these three patients with *ATP1A3*-related disorders may illustrate that the symptoms associated with the FIPWE phenotype^{1,4} are encompassed within a broader phenotypic continuum of *ATP1A3* gene mutations. Assigning overly exact phenotype-genotype relationships could promote inaccurate

interpretation of the overlapping phenotypes within the broad phenotypic spectrum of *ATP1A3*-related disorders. Furthermore, clinicians need to be aware that laboratories may report specific mutations differently, and therefore we advise that publications on the association between a new acronym and a specific gene mutation also mention the reference sequence accession number (NM_number). This could alert clinicians that their previous patients with similar phenotypes could have been reported differently by another genetic reference sequencing, in the past.

Until now, it remains unclear whether, and if so, which molecular mechanisms underlie these disease presentations. In patients with *ATP1A3* gene mutations, we hope that future insight in the spectrum of genotype-phenotype relationships will add to our understanding of the pathogenesis and to potential therapeutic strategies to prevent these devastating attacks.

References

1. Yano ST, Silver K, Young R, et al. Fever-induced paroxysmal weakness and encephalopathy, a new phenotype of *ATP1A3* mutation. *Pediatr Neurol.* 2017;73:101–105.
2. Rosewich H, Sweney MT, DeBrosse S, et al. Research conference summary from the 2014 international task force on *ATP1A3*-related disorders. *Neurol Genet.* 2017;3:e139.
3. Rosewich H, Weise D, Ohlenbusch A, Gartner J, Brockmann K. Phenotypic overlap of alternating hemiplegia of childhood and CAPOS syndrome. *Neurology.* 2014;83:861–863.
4. Nakamura Y, Hattori A, Nakashima M, et al. A *de novo* p.Arg756Cys mutation in *ATP1A3* causes a distinct phenotype with prolonged weakness and encephalopathy triggered by fever. *Brain Dev.* 2017. doi:10.1016/j.braindev.2017.09.010.

Deborah A. Sival

Department of Paediatrics, Paediatric Neurology,
Beatrix Children's Hospital, University Medical Centre Groningen,
University of Groningen, Groningen, The Netherlands
E-mail address: d.a.sival@umcg.nl

Fleur Vansenne

Annemieke H. Van der Hout

Department of Genetics, University Medical Centre Groningen,
University of Groningen, Groningen, The Netherlands

Marina A.J. Tijssen

*Department of Neurology, University Medical Centre Groningen,
University of Groningen, Groningen,
The Netherlands*

Tom J. de Koning

*Department of Paediatrics, Department of Genetics,
University Medical Centre Groningen, University of Groningen,
Groningen, The Netherlands*